

VEGETABLE PRODUCTION (BRUMDEC) - REVIEW & PROPOSED SHORT-TERM ADAPTIVE PRODUCTION ORIENTED RESEARCH PROGRAMME

IICA/JAMAICA
Miscellaneous Publication #320
ISSN-0534-5391





VEGETABLE PRODUCTION (BRUMDEC) - REVIEW AND PROPOSED SHORT-TERM ADAPTIVE PRODUCTION ORIENTED RESEARCH PROGRAMME

111111111111

by

Charles Percy Kennard Vegetable Production Specialist IICA/Jamaica

The second second

VEGETABLE PRODUCTION (BRUMDEC) - REVIEW AND PROPOSED SHORT TERM ADAPTIVE PRODUCTION ORIENTED RESEARCH PROGRAMME

Introduction

The Black River Upper Morass Development Company Limited (BRUMDEC), a wholly owned subsidiary of the Jamaica National Investment Co. Ltd., (JNIC), is the legally authorized executing agent of the Black River Morass Reclamation Project. The project is concerned with the development of approximately 11,000 acres of land in the Elim and Barton Isle area of St. Elizabeth. The Company plans to cultivate onions, peas and beans, vegetables, rice, cotton, pineapple, cassava, mango, citrus, coffee, plantains, and peanuts.

The general objectives of the project provide for the initiation of research and development for introducing improved agricultural production techniques, and to provide technical assistance to the farmers in the area. In order to achieve this objective, BRUMDEC has signed an agreement with the Inter-American Institute for Cooperation on Agriculture (IICA) for the provision of five (5) Consultants in the areas of:

Rice Production
Grain Production
Vegetable Crop Development
Irrigation and Drainage Management
Farm Management

With respect to vegetable crop development, the consultant is required to prepare a programme of adaptive production oriented research aimed at solving the problems which obtain in vegetable production in the project area. Problems to be addressed include appropriateness of vegetable crops and varieties within the context of:

- (i) Jamaican consumer acceptance
- (ii) Soil/Water considerations

: the state of the s

.

.

- (iii) Seasonality of production
- (iv) Culture and harvesting characteristics
- (v) Weed, pest and disease associations
- (vi) Seed production
- (vii) Fertilizer regimes
- (viii) Crop density and yields

2. Background

In order to develop the programme to achieve the desired objectives, it was necessary to review all the reports and recommendations pertaining to the project, to become acquainted with work done in vegetable production in Jamaica and in other countries with similar conditions as existing in the project. Discussions were held with the other consultants, BRUMDEC personnel, Ministry of Agriculture staff and other knowledgeable persons including farmers who have been involved in vegetable production.

2.1. Location

The Black River Morasses are divided into the Upper Morass and the Lower Morass. They are located in the parish of St. Elizabeth in the western part of Jamaica near the south coast. The Upper Morass which is the area being developed by the project consists of approximately 11,000 acres (4,451 hectares) of agricultural land. It is situated between Maggotty in the north and Lacovia in the south.

2.2. Climatic Conditions

Climatic conditions as it pertains to vegetable production is important in determining times of planting and harvesting, irrigation cycles, plant protection methods etc. Unfortunately, this type of information for the project is almost non-existent and figures from other locations have had to be taken as indications of what is likely to occur within the project

A second

Burning Administration

The strong of the suggestion o

 $\mathcal{L}_{\mathbf{r}}$

61

en tradición de la companya de la co

the control of the co

2.2.1. Rainfall

The table below shows the average monthly precipitation of the Santa Cruz area for the years 1931-1960.

1		1
ionth	Inches	MM
anuary	2.9	73.66
ebruary	2.9	73.66
larch	4.2	106.68
pril	8.6	218.44
iay	11.1	281.94
iune	5.9	149.86
July	5.2	132.06
wgust	9.4	238.76
September	8.8	223.52
ctober	12.7	322.58
lovem ber	6.9	175.26
ecember	2.9	73.66

*Source: Statistical Yearbook of Jamaica - 1979

The figures indicate a bimodal pattern with the rains falling in April and May, and the main rainy season starts in September and goes on into November. A marked dry season occurs from the end of November. through the end of March. However, the pattern is known to vary significantly from year to year.

2.2.2. Temperature

Temperature records for the project are not available as are the records for the areas in close proximity. The meteorological records for Jamaica as a whole indicate that in spite of elevation, the warmest months are July and August and the coolest January and February. Near sea leavels the maximum, minimum and daily temperatures are 86°F (30°C), 70°F (21°C), and 78°F (25.6°C) respectively.

1. 144

•

 $oldsymbol{eta}_{i}^{oldsymbol{a}} = oldsymbol{eta}_{i}^{oldsymbol{a}} oldsym$

en de la companya de la co

and a second to the contract of the contract o

The first part of the second section

2.2.3. Humidity

The comments made about the records of temperature for the project also apply to the humidity. The table below shows the range of relative humidity for the Kingston area. The figures indicate that all through the year during the evenings, the relative humidity is over 80%. It is lowest during the day during the months of December to March which more or less coincides with the dry season.

Range of Mean Relative Humidity (1969 - 1978)

Hour	January	February	March	April	May	June
7 a.m.	86-80	86-78	85-75	81-75	84-76	84-73
3 p.m.	66-59	66-59	71-60	71-63	72-63	73-60
Hour			_			
nour	July	August	<u>September</u>	<u>October</u>	November	<u>December</u>
7 a.m.	82-63	August 84 /6	September 88-81	90-82	87-79	86-77

^{*} Source: Statistical Yearbook of Jamaica, 1979

2.3. Major Soil Types in the Project Area

2.3.1. Morass Peat Soil - 152

The Morass Peat Soil - 152 is the most wide-spread soil in the project occupying approximately 2,900 acres. It is derived under swampy conditions from the sawgrass <u>Cladium jamaicance</u> and other related sedges. It is reported to be similar to sawgrass peat found in the Negril swamps, Everglades of Florida and many other parts of the tropics. Depending on the level of the water-table, solid black peat forms at mat on the surface which overlies rotting vegetation mixed with water in various forms of decomposition, to depths varying from one foot (0.3m) to over seven feet (2.10 m), until bluish grey clay and sandy clay is encountered (4,7).

The Morass Peat Soil has a slightly acid reaction. The potash, phosphate and magnesium content of the soil appear to be medium and adequate for most crops. The analytical data indicate a very high calcium content (3,162 to 10,714 ppm Ca within the top 6 inches (2.4 cm).

and the control of th

The second secon

the control of the second seco

. $m{\phi}$

It is felt that the high calcium content is likely to contribute to manganese, iron and zinc deficiencies (4).

2.3.2. Four Paths Series of Soils - Four Paths Clay 203 and Four Paths Loam - 204

The Four Paths Series of soils are the second most widespread soils after the Morass Peat Soil. This series together with the Linstead Clay Loam - 61 and other old alluvial soils are found in large areas of the parish of St. Elizabeth. They are generally acid, deeply weathered, strongly leached and infertile in most parts.

- 2.3.2.1. The Four Paths Clay 203 has a good depth of friable strongly structured top soil in the project area, mottling does not begin to appear until 2-3 feet (0.6-0.9 m). The zinc, copper and boron content of this soil is considered to be adequate and manganese high (15).
- 2.3.2.2. The Four Paths Sandy Loam 204, because of the fact that the highly mottled sub-soil is barely covered by a layer of sand or gravel is considered as the most infertile soil of the project. Drainage through the soil is poor as is it's moisture retention capacity (4).

2.3.3. Cashew Clay Loam - 151

The Cashew Clay Loam appears to be derived from alluvium and underlies the Morass Peat in most of the Morass. The soil is an acid heavy clay and may be regarded as being a moderate fertility. Phosphate and potash are both very low whilst calcium and magnesium are considered to be adequate (4).

2.3.4. Wallens Clay - 9

Wallens Clay in the project area is derived from yellow alluvial material. The external drainage is slow and internal drainage regarded as adequate. The top soil is neutral to slightly alkaline with the alkalinity and calcium content increasing with depth. Phosphate is low in some areas and adequate in others where phosphate was previously applied to the crop. Potash is generally low and magnesium adequate (1,4).

.

:

(x,y) = (x,y) + (x,y

• · ·

- 2.3.5. Minor Soils: Linstead Clay Loam 61
 Cave Valley Clay 116
- 2.3.5.1. <u>Linstead Clay</u> Linstead Clay 61 is an acid inland basin. Its properties include poor drainage, high acidity and low phosphate and potash. Manganese and copper are considered to be adequate and zinc and boron low. (4, 15).
- 2.3.5.2. Cave Valley Clay 116 is alkaline in reaction with adequate amounts of calcium and manganese, and phosphate and potash being low.

2.4. Cropping Pattern for Major Soils within the Project

2.4.1. Morass Peat Soil - 152

Information supplied by the staff of the Ministry of Agriculture in St. Elizabeth indicates that certain vegetables were successfully cultivated on the Morass Peat Soil within the project during the period 1974-1977 (12). Onions and irish potatoes have been reported to have been grown with limited success in the Negril area (4). In the Everglades peat in Florida which is considered to be similar to the Morass Peat, tomatoes, carrots and particularly the leafy vegetables are being successfully grown (9).

In the utilization of this soil for vegetable production, it is recommended that the peat should be ploughed to a depth of 8 inches (3.1 cm). It is also recommended that ripping should be done to take out weed roots and tubers, land preparation should be done with small tractors and planting done on beds in the flat (3).

2.4.2. Four Paths Series of Soils

2.4.2.1. The Four Paths Clay - 203 is considered to be suitable for a wide range of acid loving crops, including sugar cane, citrus, pineapple, food crops, pasture and tobacco which have been successfully grown in other parts of Jamaica. Within the project, this soil is being utilized for the cultivation of sugar-cane, peanuts, red peas, cassava and for pasture (4).

en de la composition La seconda de la composition de la comp ing the contract of the contra

A construction of the second o and the control of th $\frac{\partial u}{\partial x} = \frac{\partial u}{\partial x} + \frac{\partial u}{\partial x} +$

All the description of the second of the sec

2.4.2.2. The Four Paths Loam - 204, within the project area is mainly under pasture. Pineapple and tobacco have been suggested as possible crops. The utilization of this soil for vegetables requires certain management practices e.g. deep cultivation to break the clay pan, graded drains, planting on raised beds, liming and addition of organic matter (13).

2.4.3. Cashew Clay Loam - 151

Rice growing has been recommended for the Cashew Clay Loam - 151 because of the poor internal and external drainage. Sugar cane with extensive irrigation and pasture with or without drainage have been suggested as alternative crops.

2.4.4. Wallens Clay - 9

The Wallens Clay - has been utilized within the project for rice and sugar. It has been suggested that if it becomes necessary to change the cropping pattern, bananas and food crops should be considered if adequate drainage can be provided.

2.4.5. Minor Soils

Of the minor soils, only the Linstead Clay Loam - 61 is considered to be suitable for vegetables. The management practices recommended for this soil when they are to be utilized for vegetables are essentially similar to that of the Four Paths Loam - 204. Within the project, this soil is being fully utilized for sugar cane and as such not available for vegetable production.

2.5. Water Management

2.5.1. Irrigation

The Harza Report (3) indicated that even during the months of high total rainfall, the distribution was often so uneven that the availability in the root zone was depleted by evapotranspiration. From this,

it was inferred that supplementary irrigation may be needed during any month of the year. It was recommended that the Class IIIt and Class IIId (peat soil) should be irrigated by a sprinkler system and the remaining Class II soils by flood, furrow or sprinkler irrigation. Mention was made of the fact that mole drains are being used in a number of countries to irrigate peat soils, and in Florida mole drains are being used in combination with open drains.

It has been suggested that two crops of short duration can be cultivated in a year if sown at the appropriate time during the spring and fall season (11). For vegetables on the peat the following was recommended.

Crop	Growth Period		
Carrot	April-June, September-November		
Cabbage	September-November		
Watermelon	September-November		
Onion (green)	April-May, September-October		
Onion (bulb)	April-June, September-November		

Because of the fact that most of the vegetable production in Jamaica has been geared to coincide with the rainy seasons, this often results in a few months of surpluses (gluts) and many months of shortages. Many of the vegetable crops which are presently only being grown during the rainy seasons can be profitably grown all year round with supplementary irrigation (14). A project of the magnitude of BRUMDEC must gear itself to take care of the periods of shortage within the local markets when prices would also be advantageous. Presently none of the irrigation facilities for the vegetables are in place either on the Morass Peat or the Four Paths Series.

2.5.2. Water Table

Subsidence of peat soils due to oxidation of the organic matter and shrinkage is regarded as the most important factor in their good management. Maintenance of high water and regular flooding is

en de la companya de la co

en de la companya de la co

and the second of the second o

recommended as measures to reduce the rate of shrinkages (1,4). The recommended depth of the water table for maximum yield of certain vegetables grown in peat soils, e.g. Morass Peat are:

Crop	Feet	Meters
Tomatoes	1.5 - 2.0	0.41 - 0.61
Beans & Peas	1.5 - 2.0	0.45 - 0.61
Cabbage	1.5 - 2.0	0.45 - 0.61
Carrots	2.0	0.61
Corn (sweet)	2.0 - 2.5	0.61 - 0.75
Lettuce	2.5 - 3.0	0.75 - 0.90
Onions	2.0 - 2.5	0.61 - 0.75

2.6. Fertilizer Practices

As vegetable production is not being done presently on any of the soils of the project, no observations could have been made on the effect of fertilizer practices on production. The recommendations of the Ministry of Agriculture for the Four Paths Loam - 204 (13) and that of Harza (3) for the Morass Peat - 152 are shown below:

2.6.1. Mineral soils e.g. Four Paths Loam - 204

- 2.6.1.1. Fruit/vegetables e.g. tomatoes, okra, egg plant, snap beans, peppers, sweet corn, cucumber etc. Five (5) cwt. of 7-14-14 applied one week before planting
- 2.6.1.2. Root vegetables e.g. carrots, beets, radish. Four (4) cwt. of 6-18-27 or 12-14-12 applied one week before planting.
- 2.6.1.3. Leafy vegetables e.g. cabbage, lettuce, cauliflower, celery. Three (3) cwt. of 7-14-14 applied prior to planting plus three (3) cwt. sulphate of ammonia applied as a side-dressing, 3 weeks after planting.
- 2.6.1.4. Onions. Six (6) cwt. of 7-14-14 or 4 cwt. of 12-24-12 applied seven (7) days before planting plus two (2) cwt. of sulphate of ammonia applied as side dressing 6-8 weeks after planting.

.

....

. .

.

e de la companya de l

2.6.2. Morass Peat Soil - 152

- 2.6.2.1. Onions, carrots, beets, etc. Muriate of potash, 900 lbs per acre per crop. Triple superphosphate, 300 lbs per acre per crop. One half of the fertilizer is to be applied as a basal application, and the remainder in three or more applications.
- 2.6.2.2. Tomatoes, cabbage and lettuce etc. Muriate of potash, 400 lbs per acre per crop. Triple superphosphate, 400 lbs per acre per crop. One half of the fertilizer as basal application and the remainder in three or more applications.

2.6.3. Micro-Nutrients

With few exceptions all the major alluvial soils including the Four Path Series are fairly well supplied with the micro-nutrients e.g. zinc, manganese, copper and boron (15). The application of micro-elements to crops grown on alluvial soils is not generally advocated. However, for the peat soils, micro-nutrients are considered to be necessary for successful production of vegetables. Hewitt (4) recommends are pre-plant amendments in the last stages of land preparation, the application of 15 lbs of copper, 10 lbs of manganese, 4 lbs of boron, and 5-7 lbs of zinc per acre. Except for zinc, a similar recommendation has been made by Harza (3). Other reports indicate that the application of copper may be lessened and may not be needed at all, whilst manganese and zinc may best be applied as foliar sprays.

2.7. Plant Protection

Under the conditions prevailing within the project area e.g. high temperatures and humidity, pests, diseases and weeds can be expected to be a serious problem in the production of vegetables. Appendix I and II list some of the more important pests and diseases of vegetable and their control. Appendix III gives the chemical control of veeds in selected vegetables.

• •

the second

....

with the second of the second

. .

1944 - 1944 - 1944 - 1944 - 1944 - 1944 - 1944 - 1944 - 1944 - 1944 - 1944 - 1944 - 1944 - 1944 - 1944 - 1944

 $(x_1, \dots, x_{n-1}, x_{n-1}, \dots, x_{n-1}) = \{ x_1, \dots, x_{n-1}, \dots, x_{n-1} \}$ to a second

3. Programme for Adaptive Production Oriented Research (Short Term) in Vegetable Production

3.1. Objectives

The programme for adaptive production oriented research in vegetable production has as its overall objective the following:

- 3.1.1. Establishing the appropriateness of vegetables and varieties for production within the project taking into consideration, seasonality of production, cultural practices, fertilizer regimes, yields, weeds, pests and diseases associations.
- 3.1.2. Developing a technological package of practices for implementing on-farm vegetable cultivation under conditions which prevail in the project.
- 3.1.3. Accumulating data on the various aspects of vegetable production and which will provide the basis for determining those areas for future research.
- 3.1.4. Training of national technicians consistent with the agronomy of vegetable production.

3.2. Strategy

The proposed strategy is as follows:

3.2.1. In order that the programme can produce quick benefits, main emphasis will be given to the selection of suitable crops and varieties of these crops for the cultivation on the major soils of BRUMDEC. The varieties being evaluated are those which have been found to be suitable for Jamaican conditions and to conditions similar to that of the project area. Current recommendations regarding cultural practices, fertilizer applications, pests, diseases and weed control will be followed.

and the second of the second o

 $\frac{1}{2} \left(\frac{1}{2} \left$

1.5

AND THE CONTRACTOR OF THE PARTY range king

Since the economic viability of BRUMDEC seems to lay to some extent on the utilization of the Morass Peat Soil - 152 for the production of high-priced vegetables, major emphasis would be placed on the evaluation studies of this soil. Selected vegetables would also be tried on the Four Paths Series - Four Paths Clay - 203 and Four Paths Loam - 204, the second most widespread soils in the project after the Morass Peat.

On the Morass Peat Soil, the vegetables to be evaluated included:

Cabbage, cauliflower, lettuce, peppers (sweet and hot), tomatoes, sweet corn, celery, string beans, egg plant, okra, onions (bulbing), onions (bunching), carrots, beets, garlic.

Those to be tested on the Four Paths Series are: Melons, cucumber, onions, carrots, beets, tomatoes.

- 3.2.2. Field experiments will be conducted on the Morass Peat Soil 152, Four Paths Clay 203 and Four Paths Loam 204 in order to establish the best N.P.K. treatments. The crops on which the studies will be done include tomatoes, carrots and cabbage which are taken as representing the fruit, root and leafy vegetables respectively. Onions will also be included.
- 3.2.3. Field studies will be carried out with the micro-elements Cu, Zn, Mn, Fe and Boron on the Morass Peat Soil. The objectives of these studies are to establish whether all these elements are needed, the rates to be applied and the method of application.

The crops which will be used in these studies are tomatoes, carrots, cabbages and onions.

.

e a company

4.5 · .

- 3.2.4. Selected vegetables e.g. tomatoes, melons and onions will be sown on the Four Paths Series, and cabbage, cauliflower, onions, carrots and tomatoes on the Worass Peat Soil during the different season. The objectives of these studies are to establish the optimum period for sowing and harvesting, to determine the irrigation needs of the crops when sown during the different seasons, to observe the incidence of pests, diseases and weeds and other problems associated with growing the various crops in and out of seasons.
- 3.2.5. The results obtained from the variety evaluation, N.P.K., micro-elements and time of sowing experiments (3.2.1 to 3.2.4) will be used to develop a package of practices for the production of vegetables on the Morass Peat and Four Paths Series of soils of the project. These would be evaluated in large scale commercial trials of one half to one acre.
- 3.2.6. The national technicians assigned to the vegetable programme, will as part of their training, be fully involved in all aspects of the programme. The commercial trials will be used as the basis for holding training programmes for farmers and other interested persons. It is expected that the national technicians would be adequately trained by this time to conduct the programme for farmers.

3.3. Methodology

3.1.1. Experimental design for varietal evaluation studies

The design of these studies will be a randomised complete block with four (4) replications. Plot size for each trial will be dependent on the crops being evaluated and the availability of planting material. On the Morass Peat soil a basal application of copper, manganese, boron and zinc will be made at 16.8 kg, 11.2 kg, 4.5 kg and 5.6 kg per hectare respectively. The fertilizer treatments for the different crops on the two (2) soil types are given below.

- **....**

And the second of the second o

75 ° 1 **N**

19th the control of the .

..

3.3.1.1. Morass Peat

Cabbage, cauliflower, lettuce, sweet corn, beans, celery:

 $P_2^0_5$ - 180 lbs/acre (202 kg/ha)

 K_2^0 - 240 lbs/acre (269 kg/ha)

Carrots, Beets:

 P_2O_5 - 200 lbs/acre (225 kg/ha)

K₂⁰ - 300 lbs/acre (336 kg/ha)

Onions:

 P_2O_5 - 120 lbs/acre (135 kg/ha)

K₂0 - 180 lbs/acre (202 kg/ha)

Tomatoes, peppers, egg plant:

 $P_2^0_5$ - 160 lbs/acre (180 kg/ha)

 K_2^0 - 240 lbs/acre (269 kg/ha)

Side dressing of nitrogen fertilizer will be made to the leafy vegetable if the need arises.

3.3.1.2. Four Paths Series

The recommendations of the Ministry of Agriculture which are as follows will be used:

Onions:

6 cwt. of 7-14-14 per acre plus 2 cwt of sulphate of ammonia as a side dressing.

Tomatoes, okra, beans:

5 cwt. of 7-14-14 fertilizer per acre

Carrots, beets:

4 cwt. of 6-18-27 or 12-14-12 per acre

Are the second of the second o

. .

.

erik kan sebia di se Finanzia

.

y early a

Cabbage, lettuce, cauliflower, celery:

3 cwt. of 7-14-14 fertilizer plus 3 cwt. of sulphate of ammonia as side dressing.

Routine spraying will be carried out with the appropriate chemicals for control of pests and diseases. Weed control will be done through the use of the recommended herbicides and also manually.

3.3.2. Experimental design for N.P.K. fertilizer experiments

The design of these experiments will be a 3x3x3 factorial arranged in block of nine (9) treatments with two (2) replications. The N.P.K. treatment will be:

On the Morass Peat Soil

N at 0, 40 and 80 kg/ha
P at 100, 150 and 200 kg/ha
K at 150, 250 and 350 kg/ha

On the Four Paths Series

N - 40, 80 and 120 kg/ha

P - 60, 120 and 180 kg/ha

K - 60, 120 and 180 kg/ha

All of the P and half of the N and K will be applied one week prior to planting. The remaining half of N and K will be applied depending on the crop - 3 to 6 weeks after seeding. On the Morass Peat a basal application of Copper and Boran will be applied at 16.8 Kg and 4.5 kg per acre respectively.

Manganese and Zinc would be applied as Foliar sprays at 0.375 Kg/ha in 30 - 40 gallons of water. Two applications of the Manganese and Zinc sprays will be made at 3 and 6 weeks.

The recommended cultural practices, pests, disease and weed control measures will be carried out.

Observations will be made on the growth of the plants and yields. Soil and plant analyses will be done at the appropriate times.

3.3.3 Design of micro-nutrients experiments

Two experiments are contemplated. The first experiment will be a randomised block designed and is intended to establish what micro-elements need to be applied to the Morass Peat Soil. The treatments will be as follows:

- (i) No micro-elements
- (ii) Complete package of micro-elements
 Cu 16.8 kg/ha (soil application)
 B 4.5 kg/ha (""")
 Mn 0.375 kg/ha (two foliar applications)
 Zn 0.375 kg/ha at 3 and 6 weeks
- (iii) Complete package minus Cu
- (iv) " " B
- (v) " " Mn
- (vi) " " Zn

A fertilizer application of 180 kg of P_2O_5 and 260 kg of K_2O per hectare will be made one week prior to

seeding to all the plots. Observation will be made on plant growth and yields.

Soil and plant analyses will be done at various stages before and after the application of the microelements. The recommended cultural practices, pest, disease and weed control measures will be adopted.

Based on the result of the first experiment, a factorial split plot experiment will be carried out. The micro-nutrients found to be necessary in the first experiments will be tested at various rates both as soil and foliar applications.

3.3.4 Time of planting experiments

The design of this experiment will be a randomised split block in which the main plots will be the variety and the sub-plots - the time of planting. The crop and varieties to be included in these experiments are:

Tomatoes - Varieties: Manalucie, Floradade

Tropic

Onions - Varieties: Texas Early Grano,

Desex, El Toro

Cabbage - Varieties: K.K. Cross, Roundup,

Hybrid YR

Carrots - Varieties: Danvers 126, Walt High-

color, Chantenay

The time of planting will be: November - December, February - March, June - July.

Other crops e.g. garlic, cauliflower, melons and lettuce will also be sown in observational plots to observe their performance during the different seasons.

The fertilizer applications will be the same as in experiment 3.3.1 as will be the cultural, pest, disease

and weed control practices. Irrigation will be provided to the crop when needed and records kept of the water requirements of the crop when sown during the different seasons. Observations will also be made of growth, yield, incidence of pests, diseases and weeds.

3.3.5 Commercial Trials

The commercial trials will be in units of one half to one acre. Records will be kept of the cost of production, yields and returns from the various crops. Based on the profitability of the different crops, recommendations will be made for large scale production at BRUMDEC and for future areas of research.

4. Requirements for the Programme

4.1 Personnel

The following personnel are required for the programme:

- One (1) agronomist as counterpart
- Two (2) technicians

Ten (10) workers initially - numbers to be adjusted as the programme develops.

4.2 Land

- Five (5) acres on the Morass Peat 152
- Two (2) acres on the Four Paths Clay 203
- Two (2) acres on the Four Paths Loam 204

4.3 Planting Materials (Seeds)

Appendix V gives the requirements of planting materials (seeds) for the programme.

4.4 Chemicals e.g. fertilizers, insecticides etc.

The requirements of chemicals for the programme are shown in Appendix VI.

. . ati tuli

: :

• ; the training of the second section is

 $(x_0, x_0, \dots, x_n) = (x_0, \dots, x_n)$

•

•• •

.

. .

--74 1

15 - 15 - 15 - 15 - 15

4.5 Equipment and other Items

The equipment required for land preparation, irrigation, pest control and other operations connected with the vegetable programme are listed in Appendix VII. The provision of these as well as those listed in Appendices V and VI are vital to the success of the programme.

REFERENCES

- 1. Grontmij. Black River Morasses Reclamation Project. 1964.
- 2. Stark, J. Soil and Land Use Survey, No. 14. Jamaica Parish of St. Elizabeth, 1963.
- 3. Harza. Black River Upper Morass Feasibility Report. 1977.
- 4. Hewitt, C. W. Report on the Detailed Soil Survey of the Upper Morass. 1978.
- 5. Statistical Yearbook of Jamaica. 1979.
- 6. Economic and Social Survey, Jamaica. 1980.
- 7. Arscott, T. G. The Soil of the Black River Upper Morass Drainage Project and their Cropping Pattern. 1981.
- 8. Land Use Committee Land Use Proposals for the Black River Upper Morass Development Co. Ltd. Project Oasis, Meylersfield-Hayne Development. 1981.
- 9. Shaw, M.E.A. and Hewitt, C.W. Report on a visit to Florida. 1979.
- 10. Half Year Report Black River Upper Morass Development Co. Ltd. 1981.
- 11. Pizarro, Humberto. Recommendations for Land Use and Irrigation Needs in the BRUMDEC Project. 1981.
- 12. McLeggan, Lloyd and M. Montague. Personal Communication. 1981.
- 13. Harrison, R.C. Fertilizer and Soil Management Schedule for Vegetable Growing Soils in Jamaica. 1978.
- 14. Vegetable Seminar. Crops and Soils Division, Ministry of Agriculture, June 10, 1974.
- 15. Wier, C.C.A. Trace Element Survey of the Major Soils in Jamaica. Undated.

• • entre de la companya . .•

SOME IMPORTANT PESTS OF VEGETABLES AND THEIR CONTROL

CROP	PESTS	CONTROL MEASURE
Crucifers (Cabbage, Cauliflower etc.)	Leafminers	Dimethoate (Rogor) or Fenitrothion
	Cabbage Looper (Trichoplusia ni)	Phosvel, Diazinon or Actellic
·	Cabbage Butterfly (Artogeis rapae)	Phosvel, Sevin, Atellic, Ambush or Diazinon
	Diamond Back Moth (Plutella xylostella)	Lannate, Dipterex, Phosvel, Actellic or Ambush
	Cabbage budworm (Hellula, phidilealis)	weekly application of Actellic or Gardona.
Tomato	Tomato fruitworm (Heliothis armigera)	Malathion, Sevin, Diazinon or Gelecron
	Tomato hornworm (Protoparce sexta)	Galecron, Diazinon or Fenitrothion
	Tomato Pinworm Keiferia lycopersicella)	Lannate or permethrin (Ambush)
	Leaf Sucking Bugs (Nezara sp.) (Phthia picta)	Malathion, Rogor 40 or Fenitrothion
	Thrips (Frankliniella sp)	Fenitrothion, Malathion or Rogor 40
Curcurbits		
Watermelon,	Aphids	Rogor 40, Diazinon or Malathion
•	Banded Cucumber Beetle (Diabrotica separata	Dipterex, fenitrothion or Malathion
•	Sucking Bugs - (Anasa sp) (Phthia picta)	Malathion, Dimethoate or Diazinon
	Melon Worm (Margaronia spp.)	Galecron or Diazinon

,			. •		
				•	
	. , .	•	•		
• • •					: `
	- ' ' '		;		
		the contraction			
	· -	Maritan II.			
	•	•			
	• •	· · · · · · · · · · · · · · · · · · ·			
		• .			
, ps 25					
•		•	·		
71 1 20		•			t
and the second of the second o					•
	· • • • • •				
	•	1 18 1 16 ¥			
er e er e maner e e e e e e e e e e e e e e e e e e		•	-		
•	(-	· · · · · · · · · · · · · · · · · · ·	•		
			- ••		
2. 12. 12. 14.			2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 -		
75 M. C. C. C.					
		•			
****			•		
	in the second of	•	•		
****	i de la companya de l La companya de la co		•		
**************************************	i de la companya de l La companya de la co		•		3
					\$1
			•		₹. **;5;1°; (7.3.1°)
					₹. "j.ÿ."
Table 1					
Table 1					
The second of th					
The second secon					
To also the second of the seco					
To also the second of the seco					
The second of th					
				. 	

	Leafminers	Dimenthoate, Fenitrothion or Malathion
Onions	Leafminers (Liriomyza spp)	Dimethoate or Fenitrothion
	Thrips (Thrips tabaci)	Malathion, Fenitrothion or Dimethoate
	Caterpillars	Malathion, Dipterex or Fenitrothion
Peppers	Whiteflies	Malathion or Dimethoate (Rogor 40)
Lettuce	Leafminers	Dimethoate or Fenitrothion
	Aphids	Malathion or Rogor 40
Carrots	Leafhoppers	Malathion, Sevin or Rogor 40
	Aphids	Malathion or Rogor 40
	Leafminers	Diazinon or Rogor 40
	Caterpillars	Diazinon, Phosvel or Dipterex
Beans	Leafminers (Liriomyza sp.)	Rogor 40, Diazinon, Fenitrothion
	Mites (Tetranychus spp.)	Kelthane, Fenitrothion or Rogor 40
• :	Caterpillars (Spodoptera sp.) (Pseudoplusia sp.) (Anomis sp.)	Fenitrothion, Diazinon, Dipterex, Phosvel
Corn (Sweet)	Armyworm (Spodoptera frugiperda)	Dipterex, Fenitrothion or Furadan applied to leaves and whorls of plant
	Aphids (Rhopalosiphum msidis)	Malathion, Fenitrothion or Rogor 40

•

\$ 15 miles

General Pests		(Xylomiges sunia) (Anieta infecta) (Prodenia ornitho- galli)	Application of Diazinon, Phosyel or Sevin to the soil. Poison bait con- taining Dipterex Diazinon or any suitable insecti- cide
•	Nematodes	(Heloidogyne spp.) Rotylenchulus reniformis)	Because of the wide host ranges of these two species of nematode, crop retation as a control measure is not generally successful. Soil fumigation with DECP, DD, Vydate or Furadan.
	Slugs and	Snails	Poison bait containing metaldehyde.

. 1. 1 ·:

SOME IMPORTANT DISEASES OF VEGETABLES AND THEIR CONTROL

CROP	DISEASES	CONTROL HEASURE
Beans	White mold (Sclerotina sclerotiorum)	Cultural. Chemical - some control is obtained with Benlate 50% wp at 1½ lbs per acre applied just prior to flowering and again 5-7 days later. Some control is also obtained with Difolaton at 3 pints per acre.
	Web blight (Rhizoctonia solani)	Good drainage. Controlled by spraying with Benlate, Daconil, Difolaton and Dithane-45 at the recom- mended rates at weekly inter- vals just prior or immediately after heavy rains.
	Powdery Mildew (Erysiphe polygoni)	Controlled by the application of Karathane or Benlate.
	Seed borne diseases - Anthracnose, Common blight, Mosaic	Use seed free of these diseases for sowing.
<u>Carrots</u>	Leaf and Stem Blight (Alternaria dauci)	Seed treatment with hot water at 50°C for 30 minutes or with a chemical seed dressing. Controlled by the application of Dithane-45 or Daconil. Spraying should commence with the first sign of disease or after plants are six weeks cld. Use of resistant or tolerant varieties e.g. Waltham Hi-color.
Crucifors (Cabbage, cauliflower	Downy mildew (Peronospora parasitica)	Controlled by application of Zineb, Dithane M45, Manzate D or Daconil
etc.)	Downy mildew (Rhizoctonia)	Control with Benlate 50% WP.
·	Bacterial Leaf Spot	Seed treatment with Thiram Spray with Maneb, Zineb, Antracol, Cupromaneb

APPENDIX II

(cont.)

	Bacterial Black Not (Xanthomonas campestris)	Treat seed in hotwater at 50°C for 25 minutes. After this treatment dust seed with fungicide e.g. Thiram or Coresan. Crop rotation and strict field sanitation. Grow resistant or tolerant varieties e.g. K.K. Cross, YR 50. Application of Agrimycin 200 ppm, Kocide or Cupravit
Cucurbits (cucumber, melons, etc.)	Downy mildew (Pseudoperonospora cubensis)	Use of resistant or tolerant varieties. Spray with Daconil plus Benlate combination, Antracol, Kocide, Cupravit, Dithane M 45 or Manzate
	Target Spot (Corynespora cassilicola)	Use of tolerant or resistant varieties. Spray with Manzate D + Benlate or Benlate alone or Zineb
	Powdery mildew (Oidium Spp.)	Protective spray of Captan or Dithane - 45
	Mosaic Virus	Use of disease free seeds. Roguing of infected plants at an early stage. Control of aphid vector in the field with appropriate insecticides
Lettuce	Brown Leaf Spot (Cerospora Longissima)	Control by the application of Benlate, Difolatan, Daconil or Dithane M45
Tomato	Light Blight (Phytopthora infestans)	Use of resistant and tolerant varieties e.g. Manalucie. Preventative spray prior to flowering, bi-weekly application of Cupravit Blue, Kocide, Dithane M45, Maneb or Daconil. Sanitation - use disease free transplants, destroy all infected plants.

.....

(cont.)

Farly blight (Alternari solani)	enitation - Seed treatment e.g. Thiram. Application of chemical sprays e.g. Kocide, Dithane, Maneb, Daconil at weekly intervals. Crop rotation.
Leaf Hould (Cladisporium sp.). Gray Leaf Spot (Stemphylium sp), Septoria leaf spot (Septoria sp.)	Application of Difolaton, Maneb, Cupromaneb, Cupra- vit, Kocide, Perenox or Daconil
Bacterial Spot (Xanthomonas Vesicatoria)	Combination spray of Kocide 494 plus Nufilm or Citcop plus Dithane M-45 or Agrimycin
Fusarium Wilt (Fusarium oxysporum) (Flycopersici)	No chemical control.
Southern Bacterial Wilt (Pseudomonas solanacearum)	The control measures recommended for these two wilt diseases include strict field sanitation, crop rotation, the planting of resistant or tolerant varieties e.g. Manalucie, Walter etc. tolerant to Fusarium Wilt. Good drainage.
Collar Rot (Sclerotium rolfsii)	Controlled by improving the nutritional status of the soil and by controlling nematodes
Tobacco/Tomato Mosaic Virus	Disease free seeds. Sanitation-avoid infested soil. Care should always be exercised to prevent contamination of hands with virus from tobacco products e.g. cigarettes. Roguing of fields.

·			

APPENDIX I (cont.)

Onion	Damping oft and Root R e.g. (Pythium Spp.) (Rhizoctonia solani)	I reatment of soil in the cursery with a fungicile conche.g. Thiram
	Onion Blast (Botrytus)	ise of tolerant varieties (g. red varieties are tolerant to disease but lown and white varieties (re generally susceptible. I oper drying and storage, (altural practices e.g. allow tops to dry proporty to fore harvesting.
Garlic	Purple blotch (Alternative porri)	Soil application of suitable fungicide e.g. Cupravit Blue paior to planting.
Pepper	Fusarium Wilt (<u>Fusarium</u> solani)	(introl measures as for timatoes
	Bacterial Wilt (Pseudomenas solanacearum)	Cood drainage. Roguing of fields. Avoid transplanting in the wet.
	Anthracnose (Collecto- trichum sp.) (Gloersporium sp.)	Seeds from unhealthy plants sould not be used for planting. the of tolerant varieties. Chemical sprays e.g. Khoide as weekly intervals.
, <u>,</u>	Bacterial Leaf Spot (Xanthomonas vesicator	the disease free seeds. Crop rotation. Application of Kocide once every two weeks can reduce infection.

	:			
,				

CHEMICAL RED CONTROL IN SELECTED VEGETABLES

CROPS	HERBICID)	MLTHOD OF APPLICATION
Cabbage	TOK 50% (0)	Applied post emergence when cabbage plants are in 2-3 true leaf stages and weeds quite young.
	Dacthal 75 %	Applied at 10 lbs/acre pre- emergence after seeding or transplanting. Soil must be moist.
	Diphenamid (Dymid)	Applied at 4 to 6 lbs/acre after seeding.
Celery	Kerosone oil or Stoddard solvent	Applied at 30-40 gallons/acre not later than one month after seeding or transplanting.
Carrots	Prometryn 50 MP (Gesagard)	Applied at 2-3 lbs/acre as a pre-emergence spray.
	Stoddard Solvent or Kerosene	Applied as a post emergence spray at the rate of 30-50 gallons per acre when the first true leaf has developed but before the carrots have reached the size of a lead pencil. Application should be made in the late afternoon when temperatures are below 85°F.
·:	Linuron 60% (Lorox, Afalon)	Applied at the rate of 2-3 lbs per acre when the crop has 2 or more true leaves. At the time of application, the temperature must be below 35°F.
Beans	Dacthal 75% W	Applied as a pre-emergence at 10 lbs/acre plus paraquat at 1 pt/acre if weeds are present.
Beets	Pyrazon (Pyramin 80)	Pre- or post-emergence applica- tion at 5 lbs/acre supplied wher soil is moist or incorporated in

APPENDIX 111

(cont.)

Cucurbits (Melon,. cucumbers)	Alanap 2h	P_2 - 2 gallons per acre immediately after seeding.
Corn	Atrazine (Gesaprim 80)	Pre-emergence at 14-14 1bs/acre
	Bladex 80 VP plus Lasso 4E	2ladex at 2-2½ lbs plus Lasso at 4-5 pints per acre in 20-30 gallons of water after.
Onions	Eptam plus Gramoxone	Eptam at 1/3 gallon plus Gramoxone at 1½ pint/acre applied as pre-plant application. Allow at least three weeks before seeding.
	Dacthal 75W	Applied at 10-12 lbs in 40-100 gallons of water per acre immediately after seeding.
	TOK 50 WP	Applied at 6-8 lbs/acre in 40-80 gallons of water per acre when the onions are at least in the 2-3 true leaf stage.
Tomato	Dymid 80 W	5-7 lbs (4 to 6 lbs active) per acre after direct seeding or transplanting or within one month of transplanting or seeding.
	Dacthal 75 H	10 lbs per acre, 4 to 6 weeks after transplanting or on direct seeded plants at 4" to 6" in height.

•

		•
. *	,	

LIST OF VARIABLE OF VEGLETCH'S RECOMMENDED FOR JAMAICA

CROT	VARIETUS	REMARKS
iomatoes	Ymalucie	
	Fioradei	
	Florida Mi-1	
	Homestead	
	Tropics	
	Walter	
	Roma	-
	Floradade	
Onions	Texas tarly Grand Granex Fi	Recommended for sowing in fall season (October end of
-	New Mexican White	November) Recommended for
	yellow Yellow	spring season
	El Toro	March - April
•	Granex Fi	· ····································
	Dessex	
	170330K	
Carrots	Danvers Half Long	
	Sparton Bonus	
Beets	Detroit Dark REJ	
Cucumbers	Ash1ey	Slicing cucumbers
	Poinsetta	1 31
	Straight Eight	11 **
	Explorer	Pickling cucumbers
	Spartan Dawn	" is
	Pickie	11
	Wisconsin Sn Ri	tt ir
•	MISCORSER OR KI	
Watermelon	Charleston Grey	
Haskmelon	Swiss Perfect	
Sweet Peppers	California Wonder	
	New Improved Californ	iia
	Vonde r	•
	Yolo Wonder	
Egg Plant	Black Beauty	
_ 	Long Purple	

			•	
		•		
•				
				i

APPENDIX IV

(cont.)

Cauliflower	Farly Paths Once Peak Snow Crown Monarch Fropical Sureheert	Fall	planting
Lettuce	Mignonetto Minetto Great Lakes	Leaf Head	Variety
Cabbage	K.K. Cross K.Y. Cross Y.R. Cross Roundup King Cole Superette Early Cross		

Source of information. Miss J. Brand, Ministry of Agriculture, Kingston

• •

	·		

PERSONAL (FLADS)

CROP	VARIFILES	QUANTITY (LBS)
Cabbage	King Cole	1
00000	Rio Verde	l
	Roundup	1
	K.K. Gross (sterme) stri	
•	Hybrid YR Cross 20	1
	Bonanza .	
	Shamrock	1
		1
,	Hybrid YR (summer 50)	1
	Danish Bald Head	1
Lettuce	Great Lakes	1/2
	Mignonette	1,
•	Mesa	<u>1</u>
	Minetto	1,
	Fulton	1.3 1.5
	Bibb	12
	Prizehead	13
	Plack-seed Simpson	1,
	72mpst:1	4
Cauliflower	Fangshan Extra Early	15
	Farmers Early #3	14
	Snow Queen	1/2
	Early Patna	15
Peppers (hot)	Long Cayenne	1 ₅
, , , ,	Scotch Bonnet	1/2
Peppers (sweet)	Early Calwonder	1 ₅
oppose (energy	Yolo Wonder	15
	Resistant Giant	
	California Wonder	12 15
	Wonder Ciant	1 ₂
	wonder Granc	3
Tomatoes	Walter	1
•	Floradel	1
	Tropic	1
	Madhead or Quickie	1
	(local variety from	
	St. Elizabeth)	
•	Manapal	1
	H 118	1
	Manalucie	ī
•	Calypso	i
•	Castlex 1025	î
	Ox-heart	i
	Floradade	i
	Roma	î
		•

	·	
4		

		(cont.)
Sweet Corr	(w.st Tennessa	5
	Sunshine State	7,
	USA - 34	ς.
	Iobelle	5
•	Golden Cup	5
	Florigold	5
Vatermelon	Fengshan #1 Seedless	1
	Sweet Baby	1
	Sugar Baby	1
•	Charleston Grey	i
	Jubilee	1
Celery	Golden Self blanching Dwarf Faltham Strain Summer	14
	Pascal	1 6
	Golden Plume	! ₄
String Beans	Provider	1
	Harvester	ι
	Sprite	1
	Extender	1
	Contender	1
	Eagle	1
	Tender Best	1
	Prince	1
	Top Crop	1
Egg Plant	Florida Market	k
	Pompano Harket	ig.
	Black Beauty	14
	Carly Prolific (Hybrid)	¥
	Black Bell	4
	Royado	15 15 18
, .	Florida Benuty	4
<i>:</i>	Long Purple	13
Cucumbers	Poinsett	13
	Cheroke	15 15 15
	Ashley	15
	Green Prolific	15
Okra	Clemson Spineless	14
•	Louisiana Green Volvet	¹ 4
•	Emerald	14

· ·	

		APPENDIX V
		(cont.)
Onions (butbing)	70005 (7 to 32	1
	Propicant will	1
	nev Mexican	1
	Granex Hybrac	1
	Yellow Crep o	1
	Red Creolo	1
	Dessex .	1
	Tropi Brown	1
	Hybrid Rio	1
•	£1 Toro	1
Onions (bunching)	Ebenezer	1
	White Portagal	1
•	Silverskin	1
	Green Bunching	1
Carrots	Waltham Higheolor	1
	Danvers 126	1
	Chantenay Royal	1
	Chantenay Red Core	1
	Nantes Cross Fi Hybrid	1
Beets	Detroit Dark Red	1
	Early Wonder	1
	Top Market	i
	King Red	ī
	Early Wonder Crossby	-
	Improved	1

::

AMPENDIX VI

THE LOSE SECTION OF SHOW THE ROOM BESILDES

Fertilizors	(V)	Amount
	7-18-2 12-14-12 fortilizer	6 cwt.
	Object (1997)	1,800 lbs
	Triple Superphosphate (45-50% P _e C _g)	2,600 lbs
	Murinte o` Potash (60% ∑)	3,200 lbs
	7.14 14 fertilizor	6 cwt.
	danganose sulphote	200 lbs
	Copper sulphate	200 lbs
	Iron chelate	10 1bs
	Zinc sulphate	30 lbs
	Borax	50 lbs
Insecticides and Fungicides	Metaldehyde Bait	50 lbs
mgreraes	Dipterex 80-90% SP	50 lbs
	Actellic	l gallon
	Dimethoate (Rogor 40, Porfekthion)	l gallon
	Fornasan 75-W	10 lbs
• :	Renlate 50 W	20 lbs
	Ambush	1 gallon
	Kocide 101 or Cupravit Blue	50 1bs
	Dithane 11 45	50 lbs
	Furadan 100	100 lbs
	•	

•		
	•	

		APPENDIX VI
		(cont.)
Harbi (k)	Francisco	l gallon
	Ferosene Oil	\$ gallons
	unothal 75 U	10 1bs
	B1/adex 50 ND	10 lbs
	Lasso 4F	2 gallons
	Tok 50 PP	10 lbs
	Dymid 80 C	10 1bs

.

.

·:

e .

·			
	•		
			1

A TO FIGURE 1997 FOR AS MESSED ONE I PMS

5 raying Laurent	tyj e	Amount
	Knopsack Gorayers (4 gir)	2
	Constant Prossure Knapsack Sprayers (4 gls)	2
	Motorized astblowers (Model Solo 410)	1
Field Equipment	Irrigation Pump 21-39 with hose	2
	Garden Fractor (15-23 LP) with ploughs, harrows, rotovators etc.	1
	Portable Irrigation Unit	1
Loboratory and Field Items	Torsion Balance Scale (model DEM 2-1)	1
	Gram Scale (Canacity 550 g 16 ozs)	1
	Graduated cylinder (100 cc)	4
	Graduated cylinder (500 cc)	4
	Soil Thermometer	2
	Plastic Bags (Place 61)	500
·:	Soil Auger	1
	Fensionmeters (Quick Oraw'' Model 2900)	2
	Measuring Tape (100 ft.)	2 .
	Tally Meters	2
•	Plastic Buckets (20 i)	5
	Respirators (Seedburo No. 413)	6
	Refrigerator (for st oring seeds)	1

ř				

	-	
		APPENDIX VII
		(cont.)
Cools	Pokys	10
	Hoes	10
	Shovels	10
	Pachetes	10
	languer	4
Miscellaneous Items	Stakes	500
	Paint (white)	l gallon
•	Paint (black)	l gallon
	Brush (Marking)	10
	Cord polythene	400 meters

' :

AGRICULTURE IN JAMAICA

Collection of papers of the Office of IICA in Jamaica

<u> 1977 -</u>	19	<u>78</u>	
No. I	-	1	Fritz Andrew Sibbles, "Basic Agricultural Information on Jamaica Internal Document of Work", January 1977
No. I	-	2	Yvonne Lake, "Agricultural Planning in Jamaica", June 1977
No. I	-	3	Aston S. Wood, Ph.D., "Agricultural Education in Jamaica", September - October 1977
No. I	-	4	Uli Locher, "The Marketing of Agricultural Produce in Jamaica", November 1977
No. I	-	5	G. Barker, A. Wahab, L. A. Bell, "Agricultural Research in Jamaica", November 1977
No. I	-	6	Irving Johnson, Marie Strachan, Joseph Johnson, "Land Settlement in Jamaica", December 1977
No. I	-	7	Government of Jamaica, "Agricultural Government Policy Papers", February 1978
No. I	-	8	Jose Emilio Araujo, 'The Communal Enterprise', February 1980
No. I	•	9	IICA and MOAJ, "Hillside Farming Technology - Intensive Short Course", Vols. I and II, March 1978
No. I	- :	10	Jose Emilio Araujo, "The Theory Behind the Community Enterprise - Seminar in Jamaica", March 1978
No. I	- 1	1	Marie Strachan, "A National Programme for the Development of Hillside Farming in Jamaica", April 1978
No. I	- 13	2	D. D. Henry, "Brief Overall Diagnosis of Hillside Farming in Jamaica", April 1978
No. I	- 13	3	Neville Farquharson, "Production and Marketing of Yams in Allsides and Christiana", May 1978

The Committee and The Committee of the C

	• • • •	77,1	
<u> 19. j. vojski i postobiloma i postobilova proto</u> <u>VII gradnik i postobilova protoprava koji se koji se postobilova</u>	:	1	
	·		
	£ -	• ••	
The transfer of the second of	<u> </u>	€ .cet	
The second of the second secon	,	a .c7	
The Proposition of the Common	t ₂ -	,	
The same of the color of the color of the same of the color of the col	7	10.00	
the state of the s	8 -	i . "	
Server the server of the serve	o "	i ut	
A transfer our was trained to consider the constant of the con	, ï ·	T Wife	
thought send the mile of the following the first of the f	; :	[.ij	
<u>nime destinations received by the ending makes the first and the ending of the ending</u>		• 1 of	
<u>Providence de la marcial de la contrada de la marcial de la contrada del contrada de la contrada del contrada de la contrada del contrada del contrada de la contrada del contrada de la contrada del contrada de la contrada del contrada d</u>	~ ;	- 3 . OM	

- No. I 14 R. C. E. McDonald, A. H. Wahah, "Fertility Assessment of Newly Terraced Hillside Soils Using the Microplot Technique the Allsides Case Study", 1978
- No. I 15 IICA IDB, "Course in Preparation and Evaluation of Agricultural Projects", Vols. I and II, November 1977
- No. I 16 Neville Farquaharson, "Production and Marketing of Dasheen in Allsides and Christiana", June 1978

1978 - 1979

- No. II 1 O. Arboleda-Sepulveda (IICA-CIDIA), "Agricultural Documentation and Information Network in Jamaica", September 1978
- No. II 2 Victor Quiroga, "National Agricultural Information System", (NAIS-Jamaica) Project Profile, September 1978
- No. II 3 Joseph Johnson, "A Review on Land Reform in Jamaica for the Period 1972 1978", September 1978
- No. II 4 Neville Farquharson, "ABC of Vegetable Farming", A
 Draft High School Textbook, Vols. I, II, III and IV,
 February 1979
- No. II 5 Jerry La Gra, "Elements of an Agricultural Marketing Strategy for Jamaica", March 1979
- No. II 6 D. D. Henry, I. E. Johnson, "Agricultural Extension Service in Jamaica", March 1979

<u>1979 - 1980</u>

- No. III 1 H. R. Stennett, "Watersheds of Jamaica and Considerations for an Ordinal Scale of their Development", July 1979
- No. III 2 IICA-MAJ, 'Hillside Farming in Jamaica', A Training Seminar, December 1978
- No. III 3 A. L. Wright, A. H. Wahab, H. Murray, "Performance of Six Varieties of Red Peas (Phaseolus vulgaris L.) on a Newly Terraced Ultisol in Jamaica", September 1979
- No. III 4 IICA/Jamaica Staff, "Agro-Socio-Economic Sample Survey of Allsides Trelawny, Jamaica", September 1979

The service was a service of the ser The state of the s The Mark to the first of the state of the st $\frac{11}{2} \left(\frac{11}{2} \left(\frac{11}{2} \left(\frac{11}{2} \right) \right) + \frac{11}{2} \left(\frac{11}{2} \left(\frac{11}{2} \right) + \frac{11}{2} \left(\frac{11}{2} \left(\frac{11}{2} \right) \right) + \frac{11}{2} \left(\frac{11}{2} \left(\frac{11}{2} \right) + \frac{11}{2} \left(\frac{11}{2} \left(\frac{11}{2} \right) \right) + \frac{11}{2} \left(\frac{11}{2} \left(\frac{11}{2} \right) + \frac{11}{2} \left(\frac{11}{2} \left(\frac{11}{2} \right) \right) + \frac{11}{2} \left(\frac{11}{2} \left(\frac{11}{2} \right) + \frac{11}{2} \left(\frac{11}{2} \right) + \frac{11}{2} \left(\frac{11}{2} \left(\frac{11}{2} \right) + \frac{11}{2} \left($ The state of the s The facility of the second of 1. 1. W. 2008 proliter with a figure way. The second of th gradiente de la majorita de jugações de la jugações de la filo de

No. III - 5 IICA-MOAJ, "An Approach to Agricultural Settlement of Hilly Lands", October 1979 No. III - 6 IICA-MOAJ, 'Tree Crops of Economic Importance to Hillside Farms in Jamaica", October 1979 No. III - 7 Canute McLean, "Production and Marketing of Peanuts", November 1979 1980 No. IV - 1 Joseph Johnson, "Production and Marketing of Red Peas in the Hilly Areas of Jamaica", January 1980 No. IV - 2 Lyn Snuffer, "Rural Women: An Annotated Caribbean Bibliography with special reference to Jamaica", January 1980 No. IV - 3 Vincent Campbell, Abdul Wahab, Howard Murray, "Response of Peanut (Arachis hypogaea L.) on a Newly Terraced Ultisol in Jamaica", January 1980 No. IV - 4 P. Aitken, A. Wahab, I. Johnson, A. Sahni, "Agro-Socio-Economic Survey - Pilot Hillside Agricultural Project 'PHILAGRIP' Southern Trelawny", February 1980 No. IV - 5 Glenys H. Barker, "Bibliography of Literature relating to Research and Development in the Agricultural Sector of Jamaica 1959 - 1979", March 1980 No. IV - 6 Milton R. Wedderburn, "Allsides Farmers' Pre-Cooperative A Socio-Economic Assessment", March 1980 No. IV - 7 Adele J. Wint, 'The Role of Women in the Development Process", April 1980 No. IV - 8 Milton R. Wedderburn, "The Co-operative Input in the Development of the Pilot Hillside Agricultural Project (PHILAGRIP)", April 1980 No. IV - 9 MOJ/IICA/CARDI, Fruit Trees Seminar - "Research & Development of Fruit Trees", June 1980 No. IV - 10 Henry Lancelot, 'Traditional Systems in Hillside Farming, Upper Trelawny, Jamaica", June 1980

 $\phi = a_1$

 $\frac{p_{1}-p_{2}}{p_{1}-p_{2}} = \frac{p_{2}-p_{3}}{p_{1}-p_{3}} = \frac{p_{2}-p_{3}}{p_{3}-p_{3}} = \frac{p_{3}-p_{3}}{p_{3}-p_{3}} = \frac{p_$

The Maria Control of the State of the State

- No. IV 11 IICA/Jamaica, "Pilot Hillside Agricultural Project", (PHILAGRIP), Project Document. Vols. I, II and III, June 1980
- No. IV 12

 A. Wahab, I. Johnson, P. Aitken, H. Murray and
 H. Stennett, "Highlights of the Pilot Hillside
 Agricultural Project at Allsides", July 1980
- No. IV 13

 I. Johnson, A. Wahab, P. Aitken, H. Payne, "Benchmark for a Project Profile for Developing a Peanut Industry in Jamaica", July 1980
- No. IV 14 P. Aitken, A. Wahab, I. Johnson, "The Allsides Post Peasant", August 1980
- No. IV 15 Norma Munguia, Percy Aitken, Abdul Wahab, Irving Johnson, "Salt Extraction by Solar Energy", A Miniproject, September 1980
- No. IV 16

 Abdul H. Wahab, Percy Aitken-Soux, Irving E. Johnson and Howard Murray, "The Allsides Project in Jamaica Developmental Potentials of Hillside Agriculture", September 1980
- No. IV 17

 P. Aitken, A. Wahab, I. Johnson, A. Sahney and N. Munguia, "Rural Women Survey", Vols. I, II and III, October 1980
- No. IV 18
 P. Aitken, I. E. Johnson, A. Wahab, "Assessment of Employment Among Small Hillside Farmers of Jamaica", November 1980
- No. IV 19 IICA/Jamaica "Pilot Hillside Agricultural Project", (PHILAGRIP), Final Project Document. October 1980.
- P. Aitken, A. Wahab, I. E. Johnson, Bo-Myeong Woo,
 "IICA Evaluation of the First Phase FSB Allsides
 Project", (Internal Document of Work), November 1980
- No. IV 21 MINAG/IICA/CARDI "Seminar on Multiple Cropping", December 1980

No. V - 1

N. Munguia, P. Aitken, A. Wahab, I. Johnson, "Smoke
Curing of Fish (as a household industry in Rural Jamaica)",
January 1981

- - and the second of the second o

 - $\frac{1}{12} \left(\frac{1}{12} + \frac{1}{12}$

No. V - 2	It's Relation to the Agricultural Sector and Considerations for its Management", January 1981
No. V - 3	D. D. Henry, J. R. Gayle, "The Culture of Grafted Pimento (as spice crop for Allsides, Jamaica)", January 1981
No. V - 4	Abdul H. Wahab, Noel Singh, "Agricultural Research in Jamaica", February 1981
No. V - 5	P. Aitken-Soux, A. H. Wahab, I. E. Johnson, "Country Level Action Plan (CLAP)", May 1981
No. V - 6	P. Aitken-Soux, A. H. Wahab, I. E. Johnson, 'Overview of Agricultural Development in Jamaica', May 1981
No. V - 7	Samuel Thompson, I. E. Johnson, P. Aitken-Soux, Abdul Wahab, "The Land Development & Utilization Act 1966", July 1981
No. V - 8	Abdul Wahab, Percy Aitken-Soux, Irving Johnson, Bo-Myeong Woo, Howard Murray, Joseph Dehaney, "The Experiences of Jamaica in the Management of Agricultural Production on Hillsides", July 1981
No. V - 9	Dave Hutton, Abdul Wahab, Howard Murray, "Yield Response of Yellow Yam (Dioscorea Cayenensis) After Disinfesting Planting Material of Pratylenchus Coffeae", July 1981
No. V - 10	Elaine Montague-Gordon, Abdul H. Wahab, Joseph Dehaney and Audrey Wright, "Performance of Eleven Varieties of Dry Beans (Phaseolus vulgaris) Over Two Successive Seasons on the Hillsides of Jamaica", August 1981
No. V - 11	Dave G. Hutton, Abdul H. Wahab, "Position Paper on Root Crops in Jamaica", August 1981
No. V - 12	Percy Aitken-Soux, Abdul H. Wahab, Irving E. Johnson, "Technical Assistance for the English Speaking Caribbean (Considerations for an IICA Strategy)" (Internal Document of Work), September 1981
No. V - 13	Bo-Myeong Woo, Abdul H. Wahab, Joseph Dehaney, "Crop Production on Hillsides using non-Bench Terracing Alternative Measures for Soil Conservation (first year's results of the Olive River Soil Conservation studies)", September 1981

Andrew Commence of the Commenc

The matter Toke 100 And The Control of the Control

and the second of the production of the second of the seco

A Company of the Comp

Sept.

No. V - 14 Abdul H. Wahab, Percy Aitken-Soux, Irving E. Johnson, Bo-Myeong Woo, Howard Murray and Joseph Dehaney, "Agricultural Production on Hillsides - the Allsides Project Case Study", September 1981 No. V - 15 D. G. Hutton, A. H. Wahab and J. Dehaney, "Investigating Critical Levels of Dry Rotting of Yellow Yam (Dioscorea Cayenensis) Planting Material, the Benefits of Disinfesting the Heads of Pratylenchus Coffeae and of After-Planting Nematicide Treatments", September 1981 No. V - 16 D. G. Hutton, A. H. Wahab, H. Murray and J. Dehaney, "Critical Levels of Dry Rotting of Yellow Yam (Dioscorea Cayenensis) Planting Material and Yield Responses After Disinfesting Heads of Pratylenchus Coffeae and After Post-Plant Nematicide Applications", September 1981 E. Ayer and J. Reyes, "Seminar on Mediterranean Fruit No. V - 17 Fly", September 30, 1981 No. V - 18 Bo-Myeong Woo, "Erosion Control Works in Korea", October 1981 No. V - 19 Irving E. Johnson and Percy Aitken-Soux, "Country Level Action Plan (CLAP)" (Third Revision - Internal Document of Work), October 1981 Humberto Pizarro, "Programme of Work to Establish Guidelines No. V - 20 for the Effective Administration, Operation and Maintenance of the Irrigation and Drainage District in the BRUMDEC Project" November 1981 No. V - 21 Humberto Pizarro, 'The Operation of the Drainage System in the Black River Upper Morass Project", November 1981 No. V - 22 Humberto Pizarro, "Recommendations for Land Use and Irrigation Needs in the BRUMDEC Project", November 1981 No. V - 23 Humberto Pizarro, "Organization, Operations and Maintenance of the Irrigation System in the BRUMDEC Project", November 1981

Humberto Pizarro, "Basic Information for Planning Water

Management in the BRUMDEC Project", November 1981

No. V - 24

. . 4.7 1 engele i de la companya de la compan

en a general transfer de la companya La companya de la co


No. VI - 1	Vivian Chin, "Rice Research and Production in the BRUMDEC Project State-of-the-Art Review, Identification of Constraints and Interim Recommendations and Budget for Establishing 405 Hectares (1,000 Acres) of Rice on the Clay Soils at BRUMDEC", January 1982
No. VI - 2 ±	Vivian Chin, "Programme of Work for the Short-Term Adaptive Production Oriented Research on Rice in the BRUMDEC Project", January 1982
No. VI - 3	Claude Grand-Pierre, "Adaptive Research for Grain Production (BRUMDEC) - (A Short Term Programme)", January 1982
No. VI - 4	Claude Grand-Pierre, "Experimental Procedures for Grain Crops Research on the BRUMDEC Project", January 1982
No. VI - 5	Charles Kennard, "Summary of the Proposed Programme of Work for Adaptive Production-Oriented Research (Short-Term) in Vegetable Production in the BRUMDEC Project", January 1982
No. VI - 6	Charles Kennard, "Vegetable Production (BRUMDEC) - Review and Proposed Short Term Adaptive Production Oriented Research Programme", January 1982

FECHA DE DEVOLUCION				
2		!		
	 			

出了. 中国领域区

13.4

, <u>1</u>, •



DOCUMENTO
MICROFILMADO
Fecha: 23 DIC 1982